

Building action repertoires: memory and learning functions of the basal ganglia

Ann M Graybiel

Massachusetts Institute of Technology, Cambridge, USA

Research on the basal ganglia suggests that they are critically involved in building up sequences of behavior into meaningful, goal-directed repertoires. Work on rodents, monkeys and humans suggests that the basal ganglia act as part of a distributed forebrain system that helps to encode such repertoires through behavioral learning, and that is engaged in the expression of such repertoires once they have been internalized. The basal ganglia also may be critical to the expression of innate behavioral routines. Experimental findings on reward-based learning suggest that neural activity in the striatum and substantia nigra, pars compacta changes during behavioral learning. New evidence also suggests extreme specificity in the neural connections interrelating the basal ganglia, cerebral cortex and thalamus. Adaptive control of behavior may centrally depend on these circuits and the evaluator–reinforcement circuits that modulate them.

Current Opinion in Neurobiology 1995, 5:733–741

Introduction

A major shift is under way in work on the basal ganglia. After years of painstaking work to build up knowledge of the basic anatomy, neurochemistry and cell physiology of the system, studies are now increasingly being directed toward functional issues and toward testing hypotheses about basal ganglia function. At the same time, there is renewed emphasis on trying to understand the functions of the basal ganglia in terms of their interactions with the neocortex and other structures as parts of distributed neural circuits. This activity is attracting many new people to work on the basal ganglia, including computational scientists (see [1,2•,3•]). There is no consensus on 'the' function of the basal ganglia, but work with techniques ranging from imaging in humans to recording and chemical inactivation of basal ganglia sites in primates and rodents points to the basal ganglia as part of a highly dynamic neural system involved in adaptive control of action, not only in the motor sphere, but also at the level of planning and cognition.

The key functional theme that will be stressed here is that the basal ganglia are involved in one or more stages of the building up, storage, decoding, retrieval and expression of behavioral action plans through collaboration with the neocortex, the thalamus and the limbic system (Fig. 1). These functions, partly subsumed under such terms as procedural or skill learning, sequence learning or habit

formation, were already emphasized explicitly by Hirsh [4], Van Den Bercken and Cools [5], Mishkin and Petrie [6], and Hikosaka and Wurtz [7] many years ago and have been suggested implicitly by studies indicating that the basal ganglia act in selecting or planning behaviors or motor acts processes dependent on plasticity, active memory, and retrieval, as well as forward planning. But what precisely do the basal ganglia contribute to these functions? What are the mechanisms involved? How do we take account of the regional specializations of the basal ganglia, whereby the component nuclei have districts primarily interconnected with motor, premotor, prefrontal or posterior association cortices? What are the functions of the several loop pathways that have been implicated in modulating basal ganglia? And why, if learning and memory are so central to basal ganglia function, have these not been emphasized, until recently, in the rich history of clinical work on the basal ganglia (see [8–11] and references therein)? These questions are at the forefront of the field.

Dorsal striatum is implicated in learning and memory in the rodent

Experimental psychologists have led the way in implicating the striatum in learning and memory functions

Abbreviations

GABA— γ -aminobutyric acid; **GPe**—globus pallidus, pars externa; **GPI**—globus pallidus, pars interna; **IPSP**—inhibitory postsynaptic potential; **M1**—primary motor cortex; **MPTP**—1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine; **PET**—positron emission tomography; **SMA**—supplementary motor area; **SNr**—substantia nigra, pars reticulata; **TAN**—tonically active neuron.

and in demonstrating dissociations between memory performance disrupted by lesions of the caudoputamen and those disturbed by lesions of the hippocampus and amygdaloid complex in rodents (e.g. [12–15] and references therein). Such work suggests that the caudoputamen is necessary for the formation of stimulus–response associations that are built up through reinforcement. It also indicates that different parts of the caudoputamen have different memory-related functions (for example, the medial part is involved in conditional reward tasks, whereas the lateral part functions in stimulus–response learning).

In two new studies that directly compare striatal-, hippocampal- and amygdala-based functions in rats performing in the Morris water maze, McDonald and White [16•] and Packard *et al.* [17•] used lesions and local amphetamine injections to depress or enhance functions in these sites and provided further evidence for a ‘cue learning’ (as opposed to hippocampal ‘place learning’) function for the caudoputamen. The amygdala contributes to both types of learning. Such studies are important as they refine the notions of multiple memory systems in the brain and emphasize that, even within the striatum, different regions may serve different aspects of mnemonic processing. One interesting interpretation of the response (cue) learning paradigm is that it reflects an egocentric navigational control system for movements in extrapersonal space, as opposed to the allocentric spatial relations system of the hippocampus (see [15,18–20]).

The dorsal striatum and the substantia nigra are implicated in reward-based learning in the monkey

Neurons in the striatum have highly context-dependent firing properties, suggesting that their response patterns have been built up through learning and memory. How is this achieved? A major emphasis of work on monkeys has been on reinforcement-based learning, encouraged by the continuing series of papers by Schultz and co-workers [21•,22•] indicating that the dopamine-containing neurons of the substantia nigra pars compacta, not only those of the ventral tegmental area, fire in relation to primary rewards and reward-conditioned stimuli. The earlier studies of Rolls *et al.* (e.g. [23]) pioneered this direction. Now, two new papers [24•,25•] have documented plasticity of neural firing in the tonically active neurons (TANs) of the striatum in monkeys learning a sensorimotor conditioning task. This population response of the TANs is maintained even when the behavioral response has become highly automatic, but is diminished if dopamine is depleted in one striatum by the injection of the neurotoxin MPTP [25•].

These studies, taken together with those of Schultz and co-workers, suggest a model [26•] in which

dopaminergic nigral neurons are recruited by novel rewarding stimuli, and long-term procedural memories are built up that are expressed under conditions of modulatory control by striatal dopamine (Fig. 1a). Whether the learning actually occurs in the striatum itself or in, for example, cortical regions projecting to the striatum, is left open by these studies, as is the question of how the changes in TANs affect the phasically active neurons of the striatum. The authors suggest, however, that as learning occurs, TAN activity becomes temporally coordinated. The authors believe that such coherence could underlie periods of enhanced plasticity in the striatal circuits during behavioral learning, so that new behavioral routines could gradually be acquired [27•]. When the synchronization is lost by virtue of dopamine depletion, a coherent ‘go signal’ for action might be lost. This loss of a synchronizing cue from the basal ganglia has been suggested by Martin *et al.* [28•] to characterize patients with Parkinson’s disease.

Evidence for basal ganglia activity in learning and memory in the human

Imaging studies designed to detect activation of the basal ganglia in subjects carrying out carefully designed behavioral tasks are now directly addressing the question of basal ganglia involvement in memory mechanisms in the humans. Many of the recent PET findings have been reviewed by Brooks [29•], who emphasizes optimization (see [30•]) and selection (see [31]) — rather than learning — functions for the striatum. However, new work is supporting earlier imaging studies (e.g. [32,33]) indicating that the human striatum may indeed be activated in relation to learning tasks, procedural tasks requiring retrieval of procedural memory stores, and imagined movement and mental rehearsal of behavior [34•,35•].

Notable among the new studies are those by Jueptner and colleagues (M Jueptner *et al.*, unpublished data; M Jueptner, CD Frith, DJ Brooks, RSJ Frackowiak, RE Passingham, unpublished data; M Jueptner, CD Frith, RSJ Frackowiak, DJ Brooks, RE Passingham, unpublished data), following up on work by Jenkins *et al.* [36•] in which subjects carried out sequential button-press tasks. Jueptner and co-workers (M Jueptner *et al.*, unpublished data; M Jueptner, CD Frith, DJ Brooks, RSJ Frackowiak, RE Passingham, unpublished data; M Jueptner, CD Frith, RSJ Frackowiak, DJ Brooks, RE Passingham, unpublished data) make a strong case that the human striatum is differentially active during the learning of new, as opposed to familiar or repetitive, sequences of button-presses. Moreover, they demonstrated that the differential activation of the striatum during such new learning occurs preferentially in the anterior striatum (head of the caudate nucleus), and that the anterior putamen along with the caudate nucleus is activated in a free selection task requiring preparation




Fig. 1. Cortical and basal ganglia circuits collaborate in learning and memory functions. **(a)** Working model of how the basal ganglia may participate in the internalization of action plans during learning by experience. Cognitive information regarding intention and representations stored in working memory is processed in the prefrontal and anterior premotor cortices, both of which project strongly to the striatum, especially the caudate nucleus. These inputs are reconfigured in the striatum in relation to evaluative inputs about the significance of internal and external states, derived from the limbic system and substantia nigra pars compacta (SNc). As a result, particular action components are selected for potential activation. Processing through pallidonigral–subthalamic circuitry leads to differential activation of selected programs, which are forwarded to the cerebral cortex and to pattern generators in the brainstem and spinal cord. Over time, this system builds up behavioral routines. Parts of the system may run instinctual programs as well. Only a subset of basal ganglia pathways are shown. Pale squares indicate striosomes in the striatum (especially prominent in caudate nucleus). STN, subthalamic nucleus. **(b)** Working model of how basal ganglia may contribute to control performance of behavioral sequences. Inputs from premotor and motor cortex carrying information about action plans and inputs from parietal cortex adding sensorimotor spatial information are processed in the striatum (putamen and parts of the caudate nucleus). As in (a), activation of selected action programs, with suppression of others, occurs through pallidonigral–subthalamic circuitry. These activated programs are sent to the cerebral cortex, where they may serve as predictive controllers, and are forwarded to pattern generators in the brainstem and spinal cord. This system interacts with the learning system shown in (a), and is seen as modifiable and contingent on evaluation signals such as those from the SNc. Only a subset of basal ganglia pathways are shown. The dark square in the striatum symbolizes matrix organization of the putamen and caudate nucleus.

and decision, but not learning. By contrast, the caudate nucleus is not differentially active, as subjects carry out pre-learned or repetitive tasks. Instead, the main activation is in more posterior striatal regions, including the putamen. This topography may account for the differences in motor learning versus motor control deficits reported in Parkinson's patients [9]. There is a remarkable concurrence between this anteroposterior topography in the human striatum and that found by Hikosaka and colleagues (S Miyachi, K Miyashita, Z Karadi, O Hikosaka, abstract, Fifth International Triennial Meeting of the International Basal Ganglia Society, Nemuno-Sato, Japan, May 1995, page 57) in brain lesion studies with macaque monkeys carrying out sequential button-press tasks. Jueptner and colleagues (M Jueptner, CD Frith, RSJ Frackowiak, DJ Brooks, RE Passingham, unpublished data) make the further

point that there is a general anteroposterior topography governing planning and executive functions in the frontal cortical–basal ganglia system as a whole. Figure 1 reflects this ordering.

The basal ganglia and planning ahead: neural representations of action sequences

A key paper on the supplementary motor area (SMA) by Tanji and Shima [37•] presents evidence that neurons in this cortical area fire selectively in relation to sequences of movements carried out in a particular order. This selectivity for the serial order of movements was not shown by neurons in the primary motor

cortex (M1) recorded under the same conditions in the same monkeys. For neuroscientists working on the basal ganglia, this is an exciting result. The SMA is a prime cortical target of the pallido-thalamo-cortical projection system.

There is growing evidence that the striatum itself is a structure critical for the representation of serial order of both learned and innate movement sequences. Early and continuing work on this issue has been carried out in rodents (see [38,39] and references therein). Evidence from lesion and recording experiments suggests that in rat 'syntactic grooming' sequences are encoded in the anterior caudoputamen. In an important new paper, Kermadi and Joseph [40•] document a strong representation of sequence information in the caudate nucleus of the macaque monkey. They demonstrate that not only movement-related activity, but also sensory (target)-related activity can be sequence specific. Interestingly, both Kermadi, Joseph and co-workers [40•,41] and Kimura [42] note that in tasks involving a sequence of movements, neurons in the caudate nucleus and the putamen can fire 100 milliseconds or more before the onset of electromyographic (EMG) activity in the prime mover muscles, a finding that stands in contrast to the relatively late firing of neurons tested with non-sequential tasks ([43]; see also [44]). Thus, the idea that the basal ganglia cannot be involved in the initiation of movement because their neurons are late-firing appears not to hold for all types of movement.

It is of the greatest interest that in neurologic disorders such as obsessive-compulsive disorder (in which perseverative sequences of behavior occur in the afflicted patients), abnormal metabolic activity is present in the anterior striatum as well as in frontal and anterior cingulate areas of the neocortex [45,46]. Indirect but consistent evidence from clinical studies of patients with Parkinson's disease also suggests that the striatum—and its dopamine-containing inputs from the substantia nigra—are important for performance of sequential behaviors and for predictive control [28•,47–52]. These clinical observations reinforce the notion that the basal ganglia and associated frontal cortical regions may allow the forward planning on which coordinated behavioral action sequences depend. It still is too early to integrate these findings with the large body of information about the ventral/limbic part of basal ganglia circuitry, but new studies suggest that this may soon be possible (see e.g. [53•,54•]).

Space-time coordination and behavioral sequence formation

The organization of behavior sequences, whether learned or innate, may critically depend on neural processing in the striatum, which not only receives massive cortical inputs relevant to sequence generation,

but is also in a position to act on the cortex and on pattern generators in the brainstem [55] (Fig. 1). But what is the nature of sequence-related neural processing in the striatum, and how does it differ from and functionally integrate with sequence-related processing in the SMA and other cortical areas? These are open issues. Two recent models [26•,27•], however, do raise timing as a critical feature of neural processing in the striatum.

First, evidence suggests that many different cortical inputs need to converge on individual striatal projection neurons to bring these spiny neurons to firing threshold [56], and, further, that cortical inputs to these neurons are re-mapped systematically into convergent or divergent modules [57,58,59•,60•]. This sets up a situation in which the striatal projection neurons could act as modifiable coherence-detection elements [27•,58,59•,60•]: the hard-wired patterns of convergence of their cortical (or other) inputs may thus represent a template for neural processing, but activity-dependent temporal coordination of the inputs may determine their spike activity and hence striatal output activity.

Second, there are several sets of striatal interneuron that could act differentially to modulate the spatiotemporal dynamics of striatal input-output processing. One set, the cholinergic interneurons, is thought to correspond to the TANs, whose firing can be modified during behavioral learning. It has been suggested that these interneurons could serve a 'motor-binding' function, bringing about new patterns of temporal coherence in striatal output networks as a result of learning [26•,27•]. Another set of striatal interneurons, GABAergic neurons expressing parvalbumin, may be particularly sensitive to cortical inputs and are thought to be electrotonically coupled [61]. These, too, could help set up dynamic, modifiable activity states in striatal neurons that could influence the temporal order of striatal outputs. Simultaneous *in vivo* recording from multiple striatal neurons [62•] and *in vitro* studies of striatal neurons (e.g. [63,64]) will be valuable in addressing these issues. Of particular interest is evidence that local, fast collateral inhibition by spiny neurons may not occur in the striatum [65•], but that neuropeptides may be intrinsic slow messengers providing local modulation (T Aosaki, abstract, Fifth International Triennial Meeting of the International Basal Ganglia Society, Nemuno-Sato, Japan, May 1995, page 2).

Pendulum swings and controversies in basal ganglia circuit analysis

Work on the basic anatomy and physiology of the basal ganglia has been anything but dull. Issues related to the circuit functions include the following: identifying the cortical areas that receive the basal ganglia output; the

'parallel loop' nature of corticobasal ganglia circuits; the direct-indirect pathway model for basal ganglia anatomy; and, finally, the functions of striatal compartments.

What cortical areas receive the output of the basal ganglia? There is a growing view that much of the frontal cortex, and even part of the temporal cortex, receives basal ganglia output either by way of the pallidum or via the substantia nigra, pars reticulata (SNr), or through both routes ([66•,67•]; PL Strick, personal communication). According to a new study by Inase and Tanji [68•], even the 'motor pallidum' projects mainly to thalamic regions that do not project to M1, but, rather, to cortical regions involved in higher order aspects of motor control than those carried out by the M1. This view, in turn, suggests that the basal ganglia may primarily act in the 'selection'/'forward planning'/'active memory' roles of these cortical areas.

To what degree are corticobasal ganglia circuits parallel loops? The circuit anatomy of the basal ganglia clearly shows that the cortex and basal ganglia are interlinked (Fig. 1), but it is still a matter of debate whether there are separate circuits ('loops') for different cortical regions (e.g. for motor, supplementary motor, or lateral prefrontal cortex) [69], or an intermixing of the circuits. One of the best examples of circuit specificity involving basal ganglia connections is the pathway related to oculomotor control (the oculomotor loop), and Kori, Kata, Miyashita and colleagues [70•–72•] have now added further evidence for selectivity in this system. However, if the M1-recipient striatopallidal path leads mainly to non-M1 cortex, as suggested by Inase and Tanji [68•], parallelism of the loops cannot be absolute. Such cross-loop interactions are also suggested by Inase *et al.* (M Inase, ST Sakai, J Tanji, abstract, Fifth International Triennial Meeting of the International Basal Ganglia Society, Nemuno-Sato, Japan, May 1995, page 38) in their report documenting considerable overlap of M1 and SMA inputs to the macaque putamen. Joel and Weiner [67•] further suggest that the parallel loop model fails to take into account open 'split' circuits in the system, by which prefrontal cortex as well as premotor cortical circuits are engaged by the basal ganglia. Finally, interactions within nuclei in the basal ganglia circuits themselves could provide critical crosslinks as well [27•,58,59•,73•,74•].

Is the direct-indirect pathway model [75,76] of basal ganglia anatomy accurate? The concept that the subthalamic nucleus controls the release functions of the 'direct' basal ganglia pathway has been enormously influential in guiding new clinical treatments for Parkinson's disease and in interpreting basal ganglia functions (see [77,78•–80•,81•] and references therein). In this model, the basal ganglia can disinhibit the thalamus by way of a 'direct pathway' from striatum to the thalamus via the globus pallidus, pars interna (GPi) or SNr. Controlling or sculpting this disinhibition is the 'indirect pathway' from striatum to globus pallidus, pars externa (GPe) to subthalamic nucleus to GPi (and then

to thalamus). Major clinical efforts have been mounted to use this information therapeutically in the treatment of Parkinson's disease: both lesions and stimulation of the subthalamic nucleus or the internal pallidum have been shown to relieve parkinsonian symptoms (see [76,81•,82] and references therein), apparently restoring a balance between the direct and indirect pathways that are lost in Parkinson's disease.

Parent and Hazrati [73•,74•] now question the existence of the indirect pathway: their anatomy in squirrel monkeys suggests that the subthalamic nucleus neurons receiving GPe input do not project to GPi, but, rather, back to GPe. They suggest that GPe should be considered as a general controller of a large set of nuclei including not only the GPi and SNr but also the thalamic reticular nucleus, a major modulator of thalamic function. The authors even suggest reserving the term 'indirect pathway' for projections from GPe to the internal pallidum (and SNr).

Anatomists are seldom at a loss to add 'new' pathways to a circuit diagram, but it is rare to find them subtracting one. If confirmed, the findings of Parent and Hazrati [73•,74•] will lead to considerable revision in thinking about phenomena ranging from clinical treatment protocols to differential regulation of neuropeptides in the basal ganglia, to current computational models based on the existence of direct and indirect pathways.

Revisions are also likely to result from other new information about the physiology and anatomy of the basal ganglia. Smith *et al.* [83•] report that the neocortical and thalamic inputs to the primate (squirrel monkey) striatum are strikingly different. First, corticostriatal terminals, but not thalamocortical terminals, can converge with nigrostriatal terminals on single spines in the primate striatum. Second, thalamic afferents to the striatum are reported to contact almost exclusively striatal neurons projecting to GPi, suggesting that the thalamus has a dominant control over the direct pathway (M Sidibé, Y Smith, *Neurosci Soc Abstr* 1994, 20:783). Tepper *et al.* [84•] report further evidence that collaterals of projection neurons in the SNr directly synapse on and inhibit dopaminergic nigral pars compacta neurons. This finding has major implications for basal ganglia functioning, including possible coordination of dopaminergic activity with activity of basal ganglia output pathways originating in the nigral complex. Findings such as these allow a glimpse of the variety and degree of specificity of the neural circuits associated with the basal ganglia.

What are the functions of striatal compartments? This question might have seemed esoteric a few years ago, but there is now dramatic evidence that striosomes, as defined histochemically by a variety of methods, are selectively vulnerable during the early stages of Huntington's disease. Hedreen and Folstein [85•] have proposed that the choreic movements of patients with Huntington's disease may partly be accounted for by this early loss. Striosomes are thought to innervate

preferentially the pars compacta of the substantia nigra, and loss of such (presumably inhibitory) input could, by changing functional activity in the nigrostriatal system, lead to the augmented movement patterns found in these patients. Interestingly, the only regions of the macaque monkey's frontal cortex so far found to project robustly to striosomes are in the posterior orbital cortex and the anterior cingulate cortex, regions that in humans have been implicated in obsessive-compulsive disorder [86•]. This finding has led to the suggestion that the anterior striosomal system may be particularly involved in the expression of such behavioral disorders and their normal behavioral counterparts [86•]. Striosomes have also been centrally implicated in reinforcement-based learning models of the basal ganglia [26•,57,87•,88•], following up an earlier hypothesis of White [89] that striosomes are part of a striatal reward system because they are preferentially linked to the limbic system. The anterior bias of this prefronto-striosomal system may contribute to the selective functions of the caudate nucleus in motor learning.

Evidence is accumulating that the matrix of the striatum is patchily organized (into matrisomes) as well, and that there are probably patterns of convergence, divergence and adjacency of corticostriatal inputs to matrisomes that set up templates for dynamically adjustable neural processing in the striatum (see [27•,58,86•]). Connolly and Burns [30•] have suggested that matrisomes correspond to state spaces in the striatum, which would fit well with the idea of matrisomes being critical to the adaptive functions of the basal ganglia [27•,30•,87•].

Conclusions

Integration of information across species, across brain regions and across levels of analysis is producing a new vigor in research on the basal ganglia. The articles reviewed here suggest that the basal ganglia are centrally concerned with the building up of adaptive behavioral action plans, and with the expression of these plans, under constraints imposed by forebrain and midbrain evaluator-reinforcement systems, through thalamocortical and brainstem executive systems.

Acknowledgements

The author thanks Glenn Holm and Catherine Cooper for their help with text processing. This work was supported by NIH Javits Award NS25529.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
 - of outstanding interest
1. Percheron G, McKenzie JS, Féger J (Eds): **The basal ganglia. IV. New ideas and data on structure and function.** In *Advances in Behavioral Biology*, vol 41. New York: Plenum; 1993.
 2. Houk JC, Davis JL, Beiser DG (Eds): *Models of Information Processing in the Basal Ganglia*. Cambridge, Massachusetts: MIT Press; 1995.
 - See annotation [3•].
 3. Kimura M, Graybiel AM (Eds): *Functions of the Cortico-Basal Ganglia Loop*. Tokyo: Springer-Verlag; 1995.
 - Three recent books [1,2•,3•] provide up-to-date views of the neurobiological and theoretical work on the basal ganglia.
 4. Hirsh R: **The hippocampus and contextual retrieval of information from memory: a theory.** *Behav Biol* 1974, 12:421-444.
 5. Van Den Bercken JHL, Cools AR: **Evidence for a role of the caudate nucleus in the sequential organization of behaviour.** *Behav Brain Res* 1982, 4:319-337.
 6. Mishkin M, Petrie HL: **Some implications for the analysis of learning and retention.** In *Neuropsychology of Memory*. Edited by Squire LR, Butters N. New York: Guilford Press; 1984:287-296.
 7. Hikosaka O, Wurtz RH: **Visual and oculomotor functions of monkey *Macaca mulatta* substantia nigra pars reticulata. 3. Memory contingent visual and saccade responses.** *J Neurophysiol* 1983, 49:1268-1284.
 8. Saint-Cyr JA, Taylor AE, Lang AE: **Procedural learning and neostriatal dysfunction in man.** *Brain* 1982, 111:941-959.
 9. Jordan N, Sagar HJ: **The role of the striatum in motor learning: dissociation between isometric motor control processes in Parkinson's disease.** *Int J Neurosci* 1994, 77:153-165.
 10. Butters N, Salmon D, Heindel WC: **Specificity of the memory deficits associated with basal ganglia dysfunction.** *Rev Neurol (Paris)* 1994, 150:580-587.
 11. Bhatia KP, Marsden CD: **The behavioural and motor consequences of focal lesions of the basal ganglia in man.** *Brain* 1994, 117:859-876.
 12. McDonald RJ, White NM: **A triple dissociation of memory systems: hippocampus, amygdala, and dorsal striatum.** *Behav Neurosci* 1993, 107:3-22.
 13. Packard MG, Hirsh R, White NM: **Differential effects of fornix and caudate nucleus lesions on two radial maze tasks: evidence for multiple memory systems.** *J Neurosci* 1989, 9:1465-1472.
 14. Reading PJ, Dunnett SB, Robbins TW: **Dissociable roles of the ventral, medial and lateral striatum on the acquisition and performance of a complex visual stimulus-response habit.** *Behav Brain Res* 1991, 45:147-161.
 15. Kesner RP, Bolland BL, Dakis M: **Memory for spatial locations, motor responses, and objects: triple dissociation among the hippocampus, caudate nucleus, and extrastriate visual cortex.** *Exp Brain Res* 1993, 93:462-470.
 16. McDonald RJ, White NM: **Parallel information processing in the water maze: evidence for independent memory systems involving dorsal striatum and hippocampus.** *Behav Neural Biol* 1994, 61:260-270.

This paper presents further evidence based on Morris water maze experiments that the dorsal striatum and hippocampus in the rat can be used concurrently for different aspects of memory.

17. Packard MG, Cahill L, McGaugh JL: **Amygdala modulation of hippocampal-dependent and caudate nucleus-dependent memory processes.** *Proc Natl Acad Sci USA* 1994, **91**:8477–8481.

In this study, the authors used injections of amphetamine to augment function in the striatum, hippocampus, and amygdala. The results of this study suggest that the striatum and hippocampus are involved in cue-based and spatial learning, respectively, whereas the amygdala can influence memory storage of both types.

18. Potegal M: **Role of the caudate nucleus in spatial orientation of rats.** *J Comp Physiol Psychol* 1969, **69**:756–764.

19. Wiener SI: **Spatial and behavioral correlates of striatal neurons in rats performing a self-initiated navigation task.** *J Neurosci* 1993, **13**:3802–3817.

20. Graziano MSA, Gross CG: **A bimodal map of space: somatosensory receptive fields in the macaque putamen with corresponding visual receptive fields.** *Exp Brain Res* 1993, **97**:96–109.

21. Schultz W: **The primate basal ganglia between the intention and outcome of action.** In *Functions of the Cortico-Basal Ganglia Loop*. Edited by Kimura M, Graybiel AM. Tokyo: Springer-Verlag; 1995:31–49.

See annotation [22*].

22. Schultz W, Apicella P, Romo R, Scarnati E: **Context-dependent activity in primate striatum reflecting past and future behavioral events.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:11–28.

This paper and [21*] review the work of Schultz and his collaborators, suggesting that there is reward-related unit activity in the substantia nigra and contingent activation of striatal neurons.

23. Rolls ET, Thorpe SJ, Maddison SP: **Responses of striatal neurons in the behaving monkey. 1. Head of the caudate nucleus.** *Behav Brain Res* 1983, **7**:179–210.

24. Aosaki T, Tsubokawa H, Ishida A, Watanabe K, Graybiel AM, Kimura M: **Responses of tonically active neurons in the primate's striatum undergo systematic changes during behavioral sensorimotor conditioning.** *J Neurosci* 1994, **14**:3969–3984.

See annotation [26**].

25. Aosaki T, Graybiel AM, Kimura M: **Effects of nigrostriatal dopamine system on acquired neural responses in the striatum of behaving monkey.** *Science* 1994, **265**:412–415.

See annotation [26**].

26. Aosaki T, Kimura M, Graybiel AM: **Temporal and spatial characteristics of tonically active neurons of the primate's striatum.** *J Neurophysiol* 1995, **73**:1234–1252.

This paper and [24**, 25**] present evidence that the tonically active neurons of the striatum undergo plastic changes in their response properties during behavioral conditioning in the monkey. The expression of these changes in response properties is greatly reduced by local depletion of dopamine in the striatum by infusion of MPTP. The tonically active neurons that undergo these plastic changes have broad distributions, in part observe compartmental boundaries, and may be cholinergic interneurons of the striatum.

27. Graybiel AM, Aosaki T, Flaherty AW, Kimura M: **The basal ganglia and adaptive motor control.** *Science* 1994, **265**:1826–1831.

This paper integrates work on the corticostriatal pathway with work on the physiology of neurons in the striatum to suggest that the tonically active neurons of the striatum carry out a space-time coordination of striatal function during learning. The authors suggest that attributes of this coordination could serve in motor 'binding'.

28. Martin KE, Phillips JG, Iansek R, Bradshaw JL: **Inaccuracy and instability of sequential movements in Parkinson's disease.** *Brain* 1994, **102**:131–140.

The authors suggest that the basal ganglia provide an internal cue for synchronization of activity leading to movement generation, and that this synchronizing cue is absent or defective in patients with Parkinson's disease.

29. Brooks DJ: **The role of the basal ganglia in motor control: contributions from PET.** *J Neurol Sci* 1995, **128**:1–13.

This paper presents a comprehensive review of recent PET studies on the activation of basal ganglia during different tasks carried out by normal subjects and patients with movement disorders.

30. Connolly CI, Burns JB: **A state-space striatal model.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:163–178.

This paper presents a model of striatal function to suggest that matrixes in the striatum correspond to individual 'state-spaces.' Once these are locally computed, they can control behavior through outputs to the pallidum and substantia nigra.

31. Mink JW, Thach WT: **Basal ganglia intrinsic circuits and their role in behavior.** *Curr Opin Neurobiol* 1993, **3**:950–957.

32. Seitz RJ, Roland PE, Bohm C, Greitz T, Stone-Elander S: **Motor learning in man: a positron emission tomographic study.** *Neuroreport* 1990, **1**:57–60.

33. Seitz RJ, Roland PE: **Learning of sequential finger movements in man: a combined kinematic and positron emission tomography (PET) study.** *Eur J Neurosci* 1992, **4**:154–165.

34. Klein D, Zatorre RJ, Milner B, Meyer E, Evans AC: **Left putaminal activation when speaking a second language: evidence from PET.** *Neuroreport* 1994, **5**:2295–2297.

This paper reports a comparison of neural activation patterns seen with PET in subjects speaking their first (native) language and speaking their second language, which they learned during early childhood, but later than they learned their first language. They report that the only site differential activation when the subjects spoke the second language as opposed to the first language was in the left putamen, and speculate that this may have to do with the demanding online processing required in using the second language.

35. Decety J, Perani D, Jeannerod M, Bettinardi V, Tadary B, Woods R, Mazziotta JC, Fazio F: **Mapping motor representations with positron emission tomography.** *Nature* 1994, **371**:600–602.

This PET study of regional cerebral blood flow documents increased activation of the basal ganglia along with other regions when subjects observed movement, and differential activation of the caudate nucleus and a number of cortical areas when subjects imagined making movements.

36. Jenkins IH, Brooks DJ, Nixon PD, Frackowiak RS, Passingham RE: **Motor sequence learning: a study with positron emission tomography.** *J Neurosci* 1994, **14**:3775–3790.

This paper presents evidence from PET measurements that the putamen is activated during sequence learning and retrieval (along with other brain regions), and compares and contrasts activity patterns in the prefrontal and premotor cortex, basal ganglia, and cerebellum as subjects learn sequential button-press movements.

37. Tanji J, Shima K: **Role for supplementary motor area cells in planning several movements ahead.** *Nature* 1994, **371**:413–416.

This paper presents electrophysiological evidence in behaving monkeys that neurons in the supplementary motor areas code sequences of movements and are critically sensitive to the order in which movements within a sequence are made. They suggest that the SMA is involved in the temporal organization of behavior and forward planning.

38. Berridge KC, Whishaw IQ: **Cortex, striatum and cerebellum: control of serial order in a grooming sequence.** *Exp Brain Res* 1992, **90**:275–290.

39. Aldridge JW, Berridge KC, Herman M, Zimmer L: **Neuronal coding of serial order: syntax of grooming in the neostriatum.** *Psychol Sci* 1993, **4**:391–395.

40. Kermadi I, Joseph JP: **Activity in the caudate nucleus of monkey during spatial sequencing.** *J Neurophysiol* 1995, **74**:911–933.

This paper presents an extensive analysis of unit activity that occurs in the caudate nucleus while macaque monkeys carry out sequences of saccadic eye and arm movements. The authors found many sequence-specific responses related to the movements and to the sensory stimuli triggering them, and suggest that the caudate nucleus participates in the construction and execution of plans for sequential motor behavior.

41. Kermadi I, Jurquet Y, Arzi M, Joseph JP: **Neural activity in the caudate nucleus of monkeys during spatial sequencing [Research note].** *Exp Brain Res* 1993, **94**:352–356.

42. Kimura M: **Behaviorally contingent property of movement-related activity in the primate putamen.** *J Neurophysiol* 1990, **63**:1277–1296.
 43. Kimura M, Graybiel AM: **Role of basal ganglia in sensory motor association learning.** In *Functions of the Cortico-Basal Ganglia Loop*. Edited by Kimura M, Graybiel AM. Tokyo: Springer-Verlag; 1995:2–17.
 44. Cheruel F, Dormont JF, Amalric M, Schmied A, Farin D: **The role of the putamen and pallidum in motor initiation in the cat. I. Timing of movement-related single cell activity.** *Exp Brain Res* 1994, **100**:250–266.
 45. Baxter LR Jr, Schwartz JM, Bergman KS, Szuba MP, Guze BH, Mazzotta JC, Alazraki A, Selin CE, Ferng HK, Phelps ME: **Caudate glucose metabolic rate changes with both drug and behavioral therapy of obsessive-compulsive disorder.** *Arch Gen Psychiatry* 1992, **49**:681–689.
 46. Swedo SE, Schapiro MB, Grady CL, Cheslow DL, Leonard HL, Kumar A, Friedland R, Rapoport SI, Rapoport JL: **Cerebral glucose metabolism in childhood-onset obsessive-compulsive disorder.** *Arch Gen Psychiatry* 1989, **46**:518–523.
 47. Agostino R, Berardelli A, Formica A, Accornero N, Manfredi M: **Sequential arm movements in patients with Parkinson's disease, Huntington's disease and dystonia.** *Brain* 1992, **115**:1481–1495.
 48. Benecke R, Rothwell JC, Dick JPR, Day BL, Marsden CD: **Disturbance of sequential movements in patients with Parkinson's disease.** *Brain* 1987, **110**:361–379.
 49. Flowers KA: **Lack of prediction in the motor behavior of parkinsonism.** *Brain* 1978, **101**:35–52.
 50. Harrington DL, Haaland KY: **Sequencing in Parkinson's disease. Abnormalities in programming and controlling movement.** *Brain* 1991, **114**:99–115.
 51. Morris RG, Downes JJ, Sahakian BJ, Evenden JL, Heald A, Robbins TW: **Planning and spatial working memory in Parkinson's disease.** *J Neurol Neurosurg Psychiatry* 1988, **51**:757–766.
 52. Ventre J, Zee DS, Papageorgiou H, Reich S: **Abnormalities of predictive saccades in hemi-Parkinson's disease.** *Brain* 1992, **115**:1147–1165.
 53. Lavin A, Grace AA: **Modulation of dorsal thalamic cell activity by the ventral pallidum: its role in the regulation of thalamocortical activity by the basal ganglia.** *Synapse* 1994, **18**:104–127.
- This paper emphasizes the potential for powerful modulation of thalamic activity by the ventral part of the pallidal complex via the thalamic reticular nucleus (cf. [73•]).
54. Stern CE, Passingham RE: **The nucleus accumbens in monkeys (Macaca fascicularis): I. The organization of behaviour.** *Behav Brain Res* 1994, **61**:9–21.
- This paper presents behavioral evidence suggesting that the ventral striatum, like the dorsal striatum, is critical to organizing sequences of behavioral action.
55. Garcia-Rill E: **The basal ganglia and the locomotor regions.** *Brain Res* 1986, **396**:47–63.
 56. Wilson CJ: **The contribution of cortical neurons to the firing patterns of striatal spiny neurons.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:29–50.
 57. Graybiel AM, Kimura M: **Adaptive neural networks in the basal ganglia.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:103–116.
 58. Parthasarathy HB, Schall JD, Graybiel AM: **Distributed but convergent ordering of striatal projections: the frontal eye field and the supplementary eye field in the monkey.** *J Neurosci* 1992, **12**:4468–4488.
 59. Flaherty AW, Graybiel AM: **Input-output organization of the sensorimotor striatum in the squirrel monkey.** *J Neurosci* 1994, **14**:599–610.
- This paper presents evidence that there is a divergent-reconvergent architecture in the cortico-striato-pallidal system of monkeys.
60. Cowan RL, Wilson CJ: **Spontaneous firing patterns and axonal projections of single corticostriatal neurons in the rat medial agranular cortex.** *J Neurophysiol* 1994, **71**:17–32.
- This paper presents an elegant analysis of the up-down state firing patterns of corticostriatal neurons and demonstrates the nature of their terminal ramifications within the striatum.
61. Kita H: **GABAergic circuits of the striatum.** In *Chemical Signalling in the Basal Ganglia: Progress in Brain Research*, vol 99. Edited by Arbuthnott GW, Emson PC. Amsterdam: Elsevier; 1993:51–72.
 62. Woodward DJ, Kirillov AB, Myre CD, Sawyer SF: **Neostriatal circuitry as a scalar memory: modeling and ensemble neuron recording.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:315–336.
- This paper presents data from multielectrode recordings suggesting that striatal neurons distributed at considerable distances from one another can fire coordinately as rats carry out sequences of movements.
63. Calabresi P, Pisani A, Mercuri NB, Bernardi G: **Postreceptor mechanisms underlying striatal long-term depression.** *J Neurosci* 1994, **14**:4871–4881.
 64. Nisenbaum ES, Xu ZC, Wilson CJ: **Contribution of a slowly inactivating potassium current to the transition to firing of neostriatal spiny projection neurons.** *J Neurophysiol* 1994, **71**:1174–1189.
 65. Jaeger D, Kita H, Wilson CJ: **Surround inhibition among projection neurons is weak or nonexistent in the rat neostriatum.** *J Neurophysiol* 1994, **72**:2555–2558.
- In a study involving a technical *tour de force*, Jaeger and colleagues use dual intracellular recordings of nearby neurons in striatal slices and antidromic activation studies *in vivo* and *in vitro* to present evidence that neighboring striatal neurons do not inhibit one another, as detected by recording of IPSPs. This evidence culminates years of study examining the possibility of local inhibitory function of the well developed local collaterals of striatal spiny neurons.
66. Middleton FA, Strick PL: **Anatomical evidence for cerebellar and basal ganglia involvement in higher cognitive functions.** *Science* 1994, **266**:458–461.
- This paper uses the retrograde transneuronal virus tracing method in the monkey to demonstrate that the basal ganglia and cerebellum both send trans-thalamic inputs to area 46 of the prefrontal cortex, a region implicated in cognitive function.
67. Joel D, Weiner I: **The organization of the basal ganglia-thalamocortical circuits: open interconnected rather than closed segregated.** *Neuroscience* 1994, **63**:363–379.
- This paper reviews anatomy of basal ganglia-cerebral cortical connections and suggests that the parallel loops model needs to be modified to include open as well as closed pathways, and introduces the concept of split circuits in this connectivity.
68. Inase M, Tanji J: **Thalamic distribution of projection neurons to the primary motor cortex relative to afferent terminal fields from the globus pallidus in the macaque monkey.** *J Comp Neurol* 1995, **353**:415–426.
- This paper demonstrates that the thalamic projection from the motor region of the pallidum primarily distributes to the motor thalamus, which does not project to the motor cortex, but instead to premotor/SMA regions.
69. Hoover JE, Strick PL: **Multiple output channels in the basal ganglia.** *Science* 1993, **259**:819–821.
 70. Kato M, Miyashita N, Hikosaka O, Matsumura M, Usui S, Kori A: **Eye movements in monkeys with local dopamine depletion in the caudate nucleus. I. Deficits in spontaneous saccades.** *J Neurosci* 1995, **15**:912–927.
- See annotation [71•].
71. Kori A, Miyashita N, Kato M, Hikosaka O, Usui S, Matsumura M: **Eye movements in monkeys with local dopamine depletion in the caudate nucleus. II. Deficits in voluntary saccades.** *J Neurosci* 1995, **15**:928–941.
- This paper and its companion [70•] demonstrate that local infusions of the neurotoxin MPTP in the macaque caudate nucleus result in selective

deficits in saccadic eye movements directed toward the contralateral side. These deficits occur when the saccades are made spontaneously and in relation to visual or memory cues.

72. Miyashita N, Hikosaka O, Kato M: **Visual hemineglect induced by unilateral striatal dopamine deficiency in monkeys.** *Neuroreport* 1995, 6:1257–1260.

This paper extends work on the effects of MPTP infusions in the macaque caudate nucleus to show that they lead to a visual hemineglect syndrome.

73. Parent A, Hazrati LN: **Functional anatomy of the basal ganglia. II. The cortico-basal ganglia-thalamo-cortical loop.** *Brain Res Rev* 1995, 20:91–127.

See annotation [74**].

74. Parent A, Hazrati LN: **Functional anatomy of the basal ganglia. II. The place of subthalamic nucleus and external pallidum in basal ganglia circuitry.** *Brain Res Rev* 1995, 20:128–154.

This paper and [73**] provide an update on basal ganglia anatomy, and challenge the model of basal ganglia pathways involving 'direct' and 'indirect' pathways.

75. Albin RL, Young AB, Penney JB: **The functional anatomy of basal ganglia disorders.** *Trends Neurosci* 1989, 12:366–375.

76. DeLong MR: **Primate models of movement disorders of basal ganglia origin.** *Trends Neurosci* 1990, 13:281–289.

77. Bergman H, Wichmann T, DeLong MR: **Reversal of experimental parkinsonism by lesions of the subthalamic nucleus.** *Science* 1990, 249:1436–1438.

78. Wichmann T, Bergman H, DeLong MR: **The primate subthalamic nucleus. I. Functional properties in intact animals.** *J Neurophysiol* 1994, 72:494–506.

See annotation [80*].

79. Bergman H, Wichmann T, Karmon B, DeLong MR: **The primate subthalamic nucleus. II. Neuronal activity in the MPTP model of parkinsonism.** *J Neurophysiol* 1994, 72:507–520.

See annotation [80*].

80. Wichmann T, Bergman H, DeLong MR: **The primate subthalamic nucleus. III. Changes in motor behavior and neuronal activity in the internal pallidum induced by subthalamic inactivation in the MPTP model of parkinsonism.** *J Neurophysiol* 1994, 72:521–530.

This paper and [78*,79*] make a three-part series presenting an extensive electrophysiological study of the subthalamic nucleus in normal monkeys and in monkeys in which MPTP has induced a primate Parkinsonian syndrome.

81. Marsden CD, Obeso JA: **The functions of the basal ganglia and the paradox of stereotaxic surgery.** *Brain* 1994, 117:877–897.

This paper presents a comprehensive overview and commentary on the roles of different parts of the fronto-basal ganglia network.

82. Limousin P, Pollak P, Benazzouz A, Hoffmann D, Le Bas JF, Broussolle E, Perret JE, Benabid AL: **Effect on parkinsonian signs and symptoms of bilateral subthalamic nucleus stimulation.** *Lancet* 1995, 345:91–95.

83. Smith Y, Bennett BD, Bolam JP, Parent A, Sadikot AF: **Synaptic relationships between dopaminergic afferents and cortical or thalamic input in the sensorimotor territory of the striatum in monkey.** *J Comp Neurol* 1994, 344:1–19.

This paper presents detailed evidence that in the monkey, corticostriatal and tyrosine-hydroxylase-containing (presumably nigrostriatal) terminals converge on single spines of spiny neurons in the striatum. By contrast, such convergence was not seen for thalamostriatal and nigrostriatal terminals. This finding strongly suggests that the dopaminergic modulation of striatal throughput is highly selective and may differentially control cortical as opposed to thalamic inputs.

84. Tepper JM, Martin LP, Anderson DR: **GABA_A receptor-mediated inhibition of rat substantia nigra dopaminergic neurons by pars reticulata projection neurons.** *J Neurosci* 1995, 15:3092–3103.

The authors present evidence from extracellular recording that there are direct inhibitory projections from SNr projection neurons to substantia nigra pars compacta dopamine-containing neurons. This direct synaptic connectivity would suggest a major new way by which afferent information to the substantia nigra could influence dopamine-containing fibers, and a route by which a feed-forward signal could be sent to the dopamine system by nigral efferent cells.

85. Hedreen JC, Folstein SE: **Early loss of neostriatal striosome neurons in Huntington's disease.** *J Neuropath Exp Neurol* 1995, 54:105–120.

This paper presents convincing evidence that the striosomal compartment undergoes early differential degeneration in the brains of patients suffering from Huntington's disease.

86. Eblen F, Graybiel AM: **Highly restricted origin of prefrontal cortical inputs to striosomes in the macaque monkey.** *J Neurosci* 1995, 15:5999–6013.

This paper presents evidence that inputs from the prefrontal cortex to the striosomal system in the macaque monkey originate in localized regions of the caudal orbito-frontal cortex and anterior part of the anterior cingulate cortex, and converge on the anterior part of the striosomal system. The cortical regions projecting to striosomes have been implicated in obsessive-compulsive disorder.

87. Houk JC, Adams JL, Barto AG: **A model of how the basal ganglia generate and use neural signals that predict reinforcement.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:249–270.

This chapter presents a model of the function of striosomes, suggesting that the striosomal module may represent the 'adaptive critic' in the actor-critic architecture of Barto (see [88*]).

88. Barto AG: **Adaptive critics and the basal ganglia.** In *Models of Information Processing in the Basal Ganglia*. Edited by Houk JC, Davis JL, Beiser DG. Cambridge, Massachusetts: MIT Press; 1995:215–232.

This chapter presents the adaptive critic model of Barto in relation to the possible functions of dopamine in modulating reinforcement learning in the basal ganglia (see also [87*]).

89. White NM: **A functional hypothesis concerning the striatal matrix and patches: mediation of S-R memory and reward.** *Life Sci* 1988, 45:7–12.

AM Graybiel, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, Building E25, Room 618, Cambridge, Massachusetts 02139, USA.
e-mail: graybiel@wccf.mit.edu